

# L'Hospital's Rule and Indeterminate Forms

Department of Mathematics and Natural Sciences  
BRAC University

---

## 1. Motivation

When evaluating limits, sometimes substitution yields undefined forms such as

$$\frac{0}{0}, \quad \frac{\infty}{\infty}, \quad 0 \cdot \infty, \quad \infty - \infty, \quad 0^0, \quad 1^\infty, \quad \infty^0.$$

These are called **indeterminate forms** since the resulting limit depends on the relative rates of vanishing or growth of the functions involved.

### Theorem

Let  $f, g$  be differentiable near  $a$  (except possibly at  $a$ ). Suppose:

1.  $f(a) = g(a) = 0$  or  $f(x), g(x) \rightarrow \infty$  as  $x \rightarrow a$ ,
2.  $g'(x) \neq 0$  near  $a$ ,
3.  $\lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$  exists (finite or  $\infty$ ).

Then

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}.$$

## 2. Indeterminate Form $\frac{0}{0}$

### Example

$$\lim_{x \rightarrow 0} \frac{\sin x}{x} \Rightarrow \frac{0}{0}.$$

Apply L'Hospital:

$$\lim_{x \rightarrow 0} \frac{\cos x}{1} = 1.$$

### Example

$$\lim_{x \rightarrow 0} \frac{e^x - 1}{x} = \lim_{x \rightarrow 0} \frac{e^x}{1} = 1.$$

### 3. Indeterminate Form $\frac{\infty}{\infty}$

Example

$$\lim_{x \rightarrow \infty} \frac{3x^2 + 4x}{5x^2 - 7} \Rightarrow \frac{\infty}{\infty}.$$

Differentiate:

$$\lim_{x \rightarrow \infty} \frac{6x + 4}{10x} = \frac{3}{5}.$$

Example

$$\lim_{x \rightarrow \infty} \frac{\ln x}{x} = \lim_{x \rightarrow \infty} \frac{1/x}{1} = 0.$$

### 4. Indeterminate Form $0 \cdot \infty$

Example

$$\lim_{x \rightarrow 0^+} x \ln x = \lim_{x \rightarrow 0^+} \frac{\ln x}{1/x}.$$

Apply L'Hospital:

$$\lim_{x \rightarrow 0^+} \frac{1/x}{-1/x^2} = \lim_{x \rightarrow 0^+} (-x) = 0.$$

### 5. Indeterminate Form $\infty - \infty$

Example

$$\lim_{x \rightarrow \infty} (\sqrt{x^2 + x} - x)$$

Multiply by conjugate:

$$\frac{x}{\sqrt{x^2 + x} + x} = \frac{1}{\sqrt{1 + 1/x} + 1} \rightarrow \frac{1}{2}.$$

### 6. Exponential Indeterminate Forms

Example

$$\lim_{x \rightarrow 0^+} x^x.$$

Let  $\ln y = x \ln x$ . Since  $x \ln x \rightarrow 0$ ,

$$\ln y \rightarrow 0 \Rightarrow y \rightarrow e^0 = 1.$$

### Example

$$\lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^x.$$

Let  $\ln y = x \ln(1 + 1/x)$ .

$$\lim_{x \rightarrow \infty} x \left( \frac{1}{x} - \frac{1}{2x^2} + \cdots \right) = 1,$$

so  $y = e$ .

### Example

$$\lim_{x \rightarrow 0^+} \left(\frac{1}{x}\right)^x.$$

$\ln y = -x \ln x \rightarrow 0 \Rightarrow y \rightarrow 1$ .

## 7. Repeated Applications

### Example

$$\lim_{x \rightarrow 0} \frac{1 - \cos x}{x^2}.$$

After two applications of L'Hospital:

$$\lim_{x \rightarrow 0} \frac{\cos x}{2} = \frac{1}{2}.$$

### Example

$$\lim_{x \rightarrow 0} \frac{e^{2x} - 1 - 2x}{x^2} = \lim_{x \rightarrow 0} \frac{2e^{2x}}{1} = 2.$$

## 8. Summary Table

Type	Example	Technique	Result
$0/0$	$\frac{\sin x}{x}$	Direct L'Hospital	1
$\infty/\infty$	$\frac{\ln x}{x}$	Direct L'Hospital	0
$0 \cdot \infty$	$x \ln x$	Rewrite as $\ln x/(1/x)$	0
$\infty - \infty$	$\sqrt{x^2 + x} - x$	Multiply by conjugate	1/2
$0^0$	$x^x$	Logarithmic form	1
$1^\infty$	$(1 + 1/x)^x$	Logarithmic form	$e$
$\infty^0$	$(1/x)^x$	Logarithmic form	1

### Note

#### Key Ideas:

- L'Hospital's Rule compares *rates of change* of numerator and denominator.
- Sometimes algebraic simplification is easier than differentiation.
- Always ensure  $g'(x) \neq 0$  and both functions are differentiable in a neighborhood of  $a$ .

### Note

When both  $f$  and  $g$  vanish (or blow up), their ratio is governed by how steeply each function changes near the point:

$$\frac{f(x)}{g(x)} \approx \frac{f'(x)}{g'(x)}.$$